

15

Statistics

Short Answer Type Questions

Q. 1 Find the mean deviation about the mean of the distribution.

| | | | | | |
|------------------|----|----|----|----|----|
| Size | 20 | 21 | 22 | 23 | 24 |
| Frequency | 6 | 4 | 5 | 1 | 4 |

Sol.

| Size | Frequency | $f_i x_i$ | $d_i = x_i - \bar{x}$ | $f_i d_i$ |
|--------------|------------------|-----------------------------|---|-----------------------------|
| 20 | 6 | 120 | 1.65 | 9.90 |
| 21 | 4 | 84 | 0.65 | 2.60 |
| 22 | 5 | 110 | 0.35 | 1.75 |
| 23 | 1 | 23 | 1.35 | 1.35 |
| 24 | 4 | 96 | 2.35 | 9.40 |
| Total | 20 | 433 | | 25 |

Now,
$$\bar{x} = \frac{\sum f_i x_i}{\sum f_i} = \frac{433}{20} = 21.65$$

$$\therefore \text{MD} = \frac{\sum f_i |x_i - \bar{x}|}{\sum f_i} = \frac{25}{20} = 1.25$$

Q. 2 Find the mean deviation about the median of the following distribution.

| | | | | | |
|---------------------------|----|----|----|----|----|
| Marks obtained | 10 | 11 | 12 | 14 | 15 |
| Number of students | 2 | 3 | 8 | 3 | 4 |

Sol.

| Marks obtained | f_i | cf | $d_i = x_i - M_e$ | $f_i d_i$ |
|-----------------------|-------------------------|------------------------|---------------------------------------|-----------------------------|
| 10 | 2 | 2 | 2 | 4 |
| 11 | 3 | 5 | 1 | 3 |
| 12 | 8 | 13 | 0 | 0 |
| 14 | 3 | 16 | 2 | 6 |
| 15 | 4 | 20 | 3 | 12 |
| Total | $\sum f_i = 20$ | | | $\sum f_i d_i = 25$ |

$$\begin{aligned} \text{Now,} \quad M_e &= \left(\frac{20+1}{2}\right)\text{th item} = \left(\frac{21}{2}\right) = 10.5\text{th item} \\ \therefore M_e &= 12 \\ \therefore \text{MD} &= \frac{\sum f_i d_i}{\sum f_i} = \frac{25}{20} = 1.25 \end{aligned}$$

Q. 3 Calculate the mean deviation about the mean of the set of first n natural numbers when n is an odd number.

Sol. Consider first natural number when n is an odd i.e., 1, 2, 3, 4, ..., n , [odd].

$$\begin{aligned} \text{Mean } \bar{x} &= \frac{1+2+3+\dots+n}{n} = \frac{n(n+1)}{2n} = \frac{n+1}{2} \\ \therefore \text{MD} &= \frac{\left|1-\frac{n+1}{2}\right| + \left|2-\frac{n+1}{2}\right| + \left|3-\frac{n+1}{2}\right| + \dots + \left|n-\frac{n+1}{2}\right|}{n} \\ &= \frac{\left|-\frac{n+1}{2}\right| + \left|2-\frac{n+1}{2}\right| + \dots + \left|\frac{n-1}{2}-\frac{n+1}{2}\right|}{n} \\ &\quad + \frac{\left|\frac{n+1}{2}-\frac{n+1}{2}\right| + \left|\frac{n+3}{2}-\frac{n+1}{2}\right| + \dots + \left|\frac{2n-2}{2}-\frac{n+1}{2}\right| + \left|n-\frac{n+1}{2}\right|}{n} \\ &= \frac{2}{n} \left[1+2+\dots+\frac{n-3}{2}+\frac{n-1}{2}\right] \left(\frac{n-1}{2}\right) \text{ terms} \\ &= \frac{2}{n} \left[\frac{\left(\frac{n-1}{2}\right)\left(\frac{n-1}{2}+1\right)}{2}\right] \left[\because \text{sum of first } n \text{ natural numbers} = \frac{n(n+1)}{2}\right] \\ &= \frac{2}{n} \cdot \frac{1}{2} \left[\left(\frac{n-1}{2}\right)\left(\frac{n+1}{2}\right)\right] = \frac{1}{n} \left(\frac{n^2-1}{4}\right) = \frac{n^2-1}{4n} \end{aligned}$$

Q. 4 Calculate the mean deviation about the mean of the set of first n natural numbers when n is an even number.

Sol. Consider first n natural number, when n is even i.e., 1, 2, 3, 4, n . [even]

$$\begin{aligned} \therefore \text{Mean } \bar{x} &= \frac{1+2+3+\dots+n}{n} = \frac{n(n+1)}{2n} = \frac{n+1}{2} \\ \text{MD} &= \frac{1}{n} \left[\left|1-\frac{n+1}{2}\right| + \left|2-\frac{n+1}{2}\right| + \left|3-\frac{n+1}{2}\right| + \left|\frac{n-2}{2}-\frac{n+1}{2}\right| + \left|\frac{n}{2}-\frac{n+1}{2}\right| \right. \\ &\quad \left. + \left|\frac{n+2}{2}-\frac{n+1}{2}\right| + \dots + \left|n-\frac{n+1}{2}\right| \right] \\ &= \frac{1}{n} \left[\left|\frac{1-n}{2}\right| + \left|\frac{3-n}{2}\right| + \left|\frac{5-n}{2}\right| + \dots + \left|\frac{-3}{2}\right| + \left|\frac{1}{2}\right| + \dots + \left|\frac{n-1}{2}\right| \right] \\ &= \frac{2}{n} \left[\frac{1}{2} + \frac{3}{2} + \dots + \frac{n-1}{2} \right] \left(\frac{n}{2}\right) \text{ terms} \\ &= \frac{1}{n} \cdot \left(\frac{n}{2}\right)^2 \quad [\because \text{sum of first } n \text{ natural numbers} = n^2] \\ &= \frac{1}{n} \cdot \frac{n^2}{4} = \frac{n}{4} \end{aligned}$$

Q. 5 Find the standard deviation of first n natural numbers.

Sol.

| | | | | | | | | |
|---------|---|---|---|----|----|-----|-----|-------|
| x_i | 1 | 2 | 3 | 4 | 5 | ... | ... | n |
| x_i^2 | 1 | 4 | 9 | 16 | 25 | ... | ... | n^2 |

Now, $\sum x_i = 1 + 2 + 3 + 4 + \dots + n = \frac{n(n+1)}{2}$

and $\sum x_i^2 = 1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$

$$\begin{aligned} \therefore \sigma &= \sqrt{\frac{\sum x_i^2}{N} - \left(\frac{\sum x_i}{N}\right)^2} \\ &= \sqrt{\frac{n(n+1)(2n+1)}{6n} - \frac{n^2(n+1)^2}{4n^2}} \\ &= \sqrt{\frac{(n+1)(2n+1)}{6} - \frac{(n+1)^2}{4}} \\ &= \sqrt{\frac{2(2n^2 + 3n + 1) - 3(n^2 + 2n + 1)}{12}} \\ &= \sqrt{\frac{4n^2 + 6n + 2 - 3n^2 - 6n - 3}{12}} \\ &= \sqrt{\frac{n^2 - 1}{12}} \end{aligned}$$

Q. 6 The mean and standard deviation of some data for the time taken to complete a test are calculated with the following results

Number of observation = 25, mean = 18.2 s, standard, deviation = 3.25 s

Further, another set of 15 observations x_1, x_2, \dots, x_{15} , also in seconds, is

now available and we have $\sum_{i=1}^{15} x_i = 279$ and $\sum_{i=1}^{15} x_i^2 = 5524$. Calculate the

standard deviation based on all 40 observations.

Sol. Given,

$$n_1 = 25, \bar{x}_1 = 18.2, \sigma_1 = 3.25,$$

$$n_2 = 15, \sum_{i=1}^{15} x_i = 279 \text{ and } \sum_{i=1}^{15} x_i^2 = 5524$$

For first set, $\sum x_i = 25 \times 18.2 = 455$

$$\therefore \sigma_1^2 = \frac{\sum x_i^2}{25} - (18.2)^2$$

$$\Rightarrow (3.25)^2 = \frac{\sum x_i^2}{25} - 331.24$$

$$\Rightarrow 10.5625 + 331.24 = \frac{\sum x_i^2}{25}$$

$$\begin{aligned} \Rightarrow \sum x_i^2 &= 25 \times (10.5625 + 331.24) \\ &= 25 \times 341.8025 \\ &= 8545.0625 \end{aligned}$$

For combined SD of the 40 observations $n = 40$,

$$\begin{aligned} \text{Now} \quad \Sigma x_j^2 &= 5524 + 8545.0625 = 14069.0625 \\ \text{and} \quad \Sigma x_j &= 455 + 279 = 734 \\ \therefore \quad \text{SD} &= \sqrt{\frac{14069.0625}{40} - \left(\frac{734}{40}\right)^2} \\ &= \sqrt{351.726 - (18.35)^2} \\ &= \sqrt{351.726 - 336.7225} \\ &= \sqrt{15.0035} = 3.87 \end{aligned}$$

Q. 7 The mean and standard deviation of a set of n_1 observations are \bar{x}_1 and s_1 , respectively while the mean and standard deviation of another set of n_2 observations are \bar{x}_2 and s_2 , respectively. Show that the standard deviation of the combined set of $(n_1 + n_2)$ observations is given by

$$\text{SD} = \sqrt{\frac{n_1(s_1)^2 + n_2(s_2)^2}{n_1 + n_2} + \frac{n_1 n_2 (\bar{x}_1 - \bar{x}_2)^2}{(n_1 + n_2)^2}}$$

Sol. Let $x_i, i = 1, 2, 3, \dots, n_1$ and $y_j, j = 1, 2, 3, \dots, n_2$

$$\therefore \quad \bar{x}_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} x_i \text{ and } \bar{x}_2 = \frac{1}{n_2} \sum_{j=1}^{n_2} y_j$$

$$\Rightarrow \quad \sigma_1^2 = \frac{1}{n_1} \sum_{i=1}^{n_1} (x_i - \bar{x}_1)^2$$

and
$$\sigma_2^2 = \frac{1}{n_2} \sum_{j=1}^{n_2} (y_j - \bar{x}_2)^2$$

Now, mean \bar{x} of the given series is given by

$$\bar{x} = \frac{1}{n_1 + n_2} \left[\sum_{i=1}^{n_1} x_i + \sum_{j=1}^{n_2} y_j \right] = \frac{n_1 \bar{x}_1 + n_2 \bar{x}_2}{n_1 + n_2}$$

The variance σ^2 of the combined series is given by

$$\sigma^2 = \frac{1}{n_1 + n_2} \left[\sum_{i=1}^{n_1} (x_i - \bar{x})^2 + \sum_{j=1}^{n_2} (y_j - \bar{x})^2 \right]$$

Now,

$$\begin{aligned} \sum_{i=1}^{n_1} (x_i - \bar{x})^2 &= \sum_{i=1}^{n_1} (x_i - \bar{x}_1 + \bar{x}_1 - \bar{x})^2 \\ &= \sum_{i=1}^{n_1} (x_i - \bar{x}_1)^2 + n_1 (\bar{x}_1 - \bar{x})^2 + 2(\bar{x}_1 - \bar{x}) \sum_{i=1}^{n_1} (x_i - \bar{x}_1) \end{aligned}$$

But
$$\sum_{i=1}^{n_1} (x_i - \bar{x}_1) = 0$$

[algebraic sum of the deviation of values of first series from their mean is zero]

Also,

$$\sum_{i=1}^{n_1} (x_i - \bar{x})^2 = n_1 s_1^2 + n_1 (\bar{x}_1 - \bar{x})^2 = n_1 s_1^2 + n_1 d_1^2$$

Where,
$$d_1 = (\bar{x}_1 - \bar{x})$$

Similarly,
$$\sum_{j=1}^{n_2} (y_j - \bar{x})^2 = \sum_{j=1}^{n_2} (y_j - \bar{x}_i + \bar{x}_i - \bar{x})^2 = n_2 s_2^2 + n_2 d_2^2$$

where,
$$d_2 = \bar{x}_2 - \bar{x}$$

Combined SD,
$$\sigma = \sqrt{\frac{[n_1(s_1^2 + d_1^2) + n_2(s_2^2 + d_2^2)]}{n_1 + n_2}}$$

where,
$$d_1 = \bar{x}_1 - \bar{x} = \bar{x}_1 - \left(\frac{n_1 \bar{x}_1 + n_2 \bar{x}_2}{n_1 + n_2} \right) = \frac{n_2(\bar{x}_1 - \bar{x}_2)}{n_1 + n_2}$$

and
$$d_2 = \bar{x}_2 - \bar{x} = \bar{x}_2 - \frac{n_1 \bar{x}_1 + n_2 \bar{x}_2}{n_1 + n_2} = \frac{n_1(\bar{x}_2 - \bar{x}_1)}{n_1 + n_2}$$

$$\therefore \sigma^2 = \frac{1}{n_1 + n_2} \left[n_1 s_1^2 + n_2 s_2^2 + \frac{n_1 n_2 (\bar{x}_1 - \bar{x}_2)^2}{(n_1 + n_2)^2} + \frac{n_2 n_1 (\bar{x}_2 - \bar{x}_1)^2}{(n_1 + n_2)^2} \right]$$

Also,
$$\sigma = \sqrt{\frac{n_1 s_1^2 + n_2 s_2^2 + \frac{n_1 n_2 (\bar{x}_1 - \bar{x}_2)^2}{(n_1 + n_2)^2}}{n_1 + n_2}}$$

Q. 8 Two sets each of 20 observations, have the same standard deviation 5. The first set has a mean 17 and the second mean 22. Determine the standard deviation of the x sets obtained by combining the given two sets.

Sol. Given, $n_1 = 20, \sigma_1 = 5, \bar{x}_1 = 17$ and $n_2 = 20, \sigma_2 = 5, \bar{x}_2 = 22$

We know that,
$$\begin{aligned} \sigma &= \sqrt{\frac{n_1 s_1^2 + n_2 s_2^2 + \frac{n_1 n_2 (\bar{x}_1 - \bar{x}_2)^2}{(n_1 + n_2)^2}}{n_1 + n_2}} \\ &= \sqrt{\frac{20 \times (5)^2 + 20 \times (5)^2 + \frac{20 \times 20 (17 - 22)^2}{(20 + 20)^2}}{20 + 20}} \\ &= \sqrt{\frac{1000}{40} + \frac{400 \times 25}{1600}} = \sqrt{25 + \frac{25}{4}} = \sqrt{\frac{125}{4}} = \sqrt{31.25} = 5.59 \end{aligned}$$

Q.9 The frequency distribution

| | | | | | | |
|----------|---|----|----|----|----|----|
| x | A | 2A | 3A | 4A | 5A | 6A |
| f | 2 | 1 | 1 | 1 | 1 | 1 |

where, A is a positive integer, has a variance of 160. Determine the value of A .

Sol.

| x | f_i | f_i x_i | f_i x_i² |
|--------------|----------------------|------------------------------------|--|
| A | 2 | 2A | 2A ² |
| 2A | 1 | 2A | 4A ² |
| 3A | 1 | 3A | 9A ² |
| 4A | 1 | 4A | 16A ² |
| 5A | 1 | 5A | 25A ² |
| 6A | 1 | 6A | 36A ² |
| Total | 7 | 22A | 92A ² |
| | $n = 7$ | $\Sigma f_i x_i = 22A$ | $\Sigma f_i x_i^2 = 92A^2$ |

$$\begin{aligned} \therefore \sigma^2 &= \frac{\sum f_i x_i^2}{n} - \left(\frac{\sum f_i x_i}{n} \right)^2 \\ \Rightarrow 160 &= \frac{92A^2}{7} - \left(\frac{22A}{7} \right)^2 \\ \Rightarrow 160 &= \frac{92A^2}{7} - \frac{484A^2}{49} \\ \Rightarrow 160 &= (644 - 484) \frac{A^2}{49} \\ \Rightarrow 160 &= \frac{160A^2}{49} \Rightarrow A^2 = 49 \\ \therefore A &= 7 \end{aligned}$$

Q. 10 For the frequency distribution

| | | | | | | |
|----------|---|---|----|----|----|---|
| x | 2 | 3 | 4 | 5 | 6 | 7 |
| f | 4 | 9 | 16 | 14 | 11 | 6 |

Find the standard distribution.

Sol.

| x_i | f_i | $d_i = x_i - 4$ | $f_i d_i$ | $f_i d_i^2$ |
|--------------|-------|-----------------|---------------------|------------------------|
| 2 | 4 | -2 | -8 | 16 |
| 3 | 9 | -1 | -9 | 9 |
| 4 | 16 | 0 | 0 | 0 |
| 5 | 14 | 1 | 14 | 14 |
| 6 | 11 | 2 | 22 | 44 |
| 7 | 6 | 3 | 18 | 54 |
| Total | 60 | | $\sum f_i d_i = 37$ | $\sum f_i d_i^2 = 137$ |

$$\begin{aligned} \therefore \text{SD} &= \sqrt{\frac{\sum f_i d_i^2}{N} - \left(\frac{\sum f_i d_i}{N} \right)^2} \\ &= \sqrt{\frac{137}{60} - \left(\frac{37}{60} \right)^2} \\ &= \sqrt{2.2833 - (0.616)^2} \\ &= \sqrt{2.2833 - 0.3794} \\ &= \sqrt{1.9037} = 1.38 \end{aligned}$$

Q. 11 There are 60 students in a class. The following is the frequency distribution of the marks obtained by the students in a test.

| | | | | | | |
|------------------|-------|-----|-------|-----------|------|-------|
| Marks | 0 | 1 | 2 | 3 | 4 | 5 |
| Frequency | $x-2$ | x | x^2 | $(x+1)^2$ | $2x$ | $x+1$ |

where, x is positive integer. Determine the mean and standard deviation of the marks

Sol. ∴ Sum of frequencies,

$$\begin{aligned}
 & x - 2 + x + x^2 + (x + 1)^2 + 2x + x + 1 = 60 \\
 \Rightarrow & 2x - 2 + x^2 + x^2 + 1 + 2x + 2x + x + 1 = 60 \\
 \Rightarrow & 2x^2 + 7x = 60 \\
 \Rightarrow & 2x^2 + 7x - 60 = 0 \\
 \Rightarrow & 2x^2 + 15x - 8x - 60 = 0 \\
 \Rightarrow & x(2x + 15) - 4(2x + 15) = 0 \\
 \Rightarrow & (2x + 15)(x - 4) = 0 \\
 \Rightarrow & x = -\frac{15}{2}, 4 \\
 \Rightarrow & x = -\frac{15}{2} \quad [\text{inadmissible}] [\because x \in I^+]
 \end{aligned}$$

| x_i | f_i | $d_i = x_i - 3$ | $f_i d_i$ | $f_i d_i^2$ |
|--------------|-------------------|-----------------|------------------------|-------------------------|
| 0 | 2 | -3 | -6 | 18 |
| 1 | 4 | -2 | -8 | 16 |
| 2 | 16 | -1 | -16 | 16 |
| $A=3$ | 25 | 0 | 0 | 0 |
| 4 | 8 | 1 | 8 | 8 |
| 5 | 5 | 2 | 10 | 20 |
| Total | $\Sigma f_i = 60$ | | $\Sigma f_i d_i = -12$ | $\Sigma f_i d_i^2 = 78$ |

$$\text{Mean} = A + \frac{\Sigma f_i d_i}{\Sigma f_i} = 3 + \left(\frac{-12}{60}\right) = 2.8$$

$$\begin{aligned}
 \sigma &= \sqrt{\frac{\Sigma f_i d_i^2}{\Sigma f_i} - \left(\frac{\Sigma f_i d_i}{\Sigma f_i}\right)^2} = \sqrt{\frac{78}{60} - \left(\frac{-12}{60}\right)^2} \\
 &= \sqrt{1.3 - 0.04} = \sqrt{1.26} = 1.12
 \end{aligned}$$

Q. 12 The mean life of a sample of 60 bulbs was 650 h and the standard deviation was 8 h. If a second sample of 80 bulbs has a mean life of 660 h and standard deviation 7 h, then find the over all standard deviation.

Sol. Here, $n_1 = 60, \bar{x}_1 = 650, s_1 = 8$ and $n_2 = 80, \bar{x}_2 = 660, s_2 = 7$

$$\begin{aligned}
 \therefore \sigma &= \sqrt{\frac{n_1 s_1^2 + n_2 s_2^2}{n_1 + n_2} + \frac{n_1 n_2 (\bar{x}_1 - \bar{x}_2)^2}{(n_1 + n_2)^2}} \\
 &= \sqrt{\frac{60 \times (8)^2 + 80 \times (7)^2}{60 + 80} + \frac{60 \times 80 (650 - 660)^2}{(60 + 80)^2}} \\
 &= \sqrt{\frac{6 \times 64 + 8 \times 49}{14} + \frac{60 \times 80 \times 100}{140 \times 140}} \\
 &= \sqrt{\frac{192 + 196}{7} + \frac{1200}{49}} = \sqrt{\frac{388}{7} + \frac{1200}{49}} \\
 &= \sqrt{\frac{2716 + 1200}{49}} = \sqrt{\frac{3916}{49}} = \frac{62.58}{7} = 8.9
 \end{aligned}$$

Q. 13 If mean and standard deviation of 100 items are 50 and 4 respectively, then find the sum of all the item and the sum of the squares of item.

Sol. Here, $\bar{x} = 50$, $n = 100$ and $\sigma = 4$

$$\begin{aligned} \therefore \quad & \frac{\Sigma x_i}{100} = 50 \\ \Rightarrow \quad & \Sigma x_i = 5000 \\ \text{and} \quad & \sigma^2 = \frac{\Sigma f_i x_i^2}{\Sigma f_i} - \left(\frac{\Sigma f_i x_i}{\Sigma f_i} \right)^2 \\ \Rightarrow \quad & (4)^2 = \frac{\Sigma f_i x_i^2}{100} - (50)^2 \\ \Rightarrow \quad & 16 = \frac{\Sigma f_i x_i^2}{100} - 2500 \\ \Rightarrow \quad & \frac{\Sigma f_i x_i^2}{100} = 16 + 2500 = 2516 \\ \therefore \quad & \Sigma f_i x_i^2 = 251600 \end{aligned}$$

Q. 14 If for distribution $\Sigma(x - 5) = 3$, $\Sigma(x - 5)^2 = 43$ and total number of item is 18. Find the mean and standard deviation.

Sol. Given,

$$n = 18, \Sigma(x - 5) = 3 \text{ and } \Sigma(x - 5)^2 = 43$$

$$\begin{aligned} \therefore \quad \text{Mean} &= A + \frac{\Sigma(x - 5)}{18} \\ &= 5 + \frac{3}{18} = 5 + 0.1666 = 5.1666 = 5.17 \end{aligned}$$

and

$$\begin{aligned} \text{SD} &= \sqrt{\frac{\Sigma(x - 5)^2}{n} - \left(\frac{\Sigma(x - 5)}{n} \right)^2} \\ &= \sqrt{\frac{43}{18} - \left(\frac{3}{18} \right)^2} \\ &= \sqrt{2.3944 - (0.166)^2} = \sqrt{2.3944 - 0.2755} = 1.59 \end{aligned}$$

Q. 15 Find the mean and variance of the frequency distribution given below.

| | | | | |
|----------|-------------------|-------------------|-------------------|--------------------|
| x | $1 \leq x \leq 3$ | $3 \leq x \leq 5$ | $5 \leq x \leq 7$ | $7 \leq x \leq 10$ |
| f | 6 | 4 | 5 | 1 |

Sol.

| x | f_i | x_i | f_ix_i | f_ix_i² |
|--------------|----------------------|----------------------|-----------------------------------|---|
| 1-3 | 6 | 2 | 12 | 24 |
| 3-5 | 4 | 4 | 16 | 64 |
| 5-7 | 5 | 6 | 30 | 180 |
| 7-10 | 1 | 8.5 | 8.5 | 72.25 |
| Total | $n = 16$ | | $\Sigma f_i x_i = 66.5$ | $\Sigma f_i x_i^2 = 340.25$ |

$$\therefore \quad \text{Mean} = \frac{\Sigma f_i x_i}{\Sigma f_i} = \frac{66.5}{16} = 4.15$$

and

$$\begin{aligned} \text{variance} &= \sigma^2 = \frac{\sum f_i x_i^2}{\sum f_i} - \left(\frac{\sum f_i x_i}{\sum f_i} \right)^2 \\ &= \frac{340.25}{16} - (4.15)^2 \\ &= 21.2656 - 17.2225 = 4.043 \end{aligned}$$

Long Answer Type Questions

Q. 16 Calculate the mean deviation about the mean for the following frequency distribution.

| | | | | | |
|-----------------------|-----|-----|------|-------|-------|
| Class interval | 0-4 | 4-8 | 8-12 | 12-16 | 16-20 |
| Frequency | 4 | 6 | 8 | 5 | 2 |

Sol.

| Class interval | f_i | x_i | $f_i x_i$ | $d_i = x_i - \bar{x} $ | $f_i d_i$ |
|-----------------------|-----------------|-------|----------------------|-------------------------|---------------------|
| 0-4 | 4 | 2 | 8 | 7.2 | 28.8 |
| 4-8 | 6 | 6 | 36 | 3.2 | 19.2 |
| 8-12 | 8 | 10 | 80 | 0.8 | 6.4 |
| 12-16 | 5 | 14 | 70 | 4.8 | 24.0 |
| 16-20 | 2 | 18 | 36 | 8.8 | 17.6 |
| Total | $\sum f_i = 25$ | | $\sum f_i x_i = 230$ | | $\sum f_i d_i = 96$ |

\therefore Mean = $\frac{\sum f_i x_i}{\sum f_i} = \frac{230}{25} = 9.2$

and mean deviation = $\frac{\sum f_i d_i}{\sum f_i} = \frac{96}{25} = 3.84$

Q. 17 Calculate the mean deviation from the median of the following data.

| | | | | | |
|-----------------------|-----|------|-------|-------|-------|
| Class interval | 0-6 | 6-12 | 12-18 | 18-24 | 24-30 |
| Frequency | 4 | 5 | 3 | 6 | 2 |

Sol.

| Class interval | f_i | x_i | cf | $d_i = x_i - m_d $ | $f_i d_i$ |
|-----------------------|----------|-------|------|---------------------|----------------------|
| 0-6 | 4 | 3 | 4 | 11 | 44 |
| 6-12 | 5 | 9 | 9 | 5 | 25 |
| 12-18 | 3 | 15 | 12 | 1 | 3 |
| 18-24 | 6 | 21 | 18 | 7 | 42 |
| 24-30 | 2 | 27 | 20 | 13 | 26 |
| Total | $N = 20$ | | | | $\sum f_i d_i = 140$ |

$\therefore \frac{N}{2} = \frac{20}{2} = 10$

So, the median class is 12-18.

$$\begin{aligned}
 \therefore \text{Median} &= l + \frac{\frac{N}{2} - cf}{f} \times i \\
 &= 12 + \frac{6}{3}(10 - 9) \\
 &= 12 + 2 = 14 \\
 \text{MD} &= \frac{\sum f_i d_i}{\sum f_i} = \frac{140}{20} = 7
 \end{aligned}$$

Q. 18 Determine the mean and standard deviation for the following distribution.

| | | | | | | | | | | | | | | | |
|------------------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|
| Marks | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 |
| Frequency | 1 | 6 | 6 | 8 | 8 | 2 | 2 | 3 | 0 | 2 | 1 | 0 | 0 | 0 | 1 |

Sol.

| Marks | f_i | $f_i x_i$ | $d_i = x_i - \bar{x}$ | $f_i d_i$ | $f_i d_i^2$ |
|--------------|-----------------|----------------------|-----------------------|---------------------|------------------------|
| 2 | 1 | 2 | 2 - 6 = -4 | -4 | 16 |
| 3 | 6 | 18 | 3 - 6 = -3 | -18 | 54 |
| 4 | 6 | 24 | 4 - 6 = -2 | -12 | 24 |
| 5 | 8 | 40 | 5 - 6 = -1 | -8 | 8 |
| 6 | 8 | 48 | 6 - 6 = 0 | 0 | 0 |
| 7 | 2 | 14 | 7 - 6 = 1 | 2 | 2 |
| 8 | 2 | 16 | 8 - 6 = 2 | 4 | 8 |
| 9 | 3 | 27 | 9 - 6 = 3 | 9 | 27 |
| 10 | 0 | 0 | 10 - 6 = 4 | 0 | 0 |
| 11 | 2 | 22 | 11 - 6 = 5 | 10 | 50 |
| 12 | 1 | 12 | 12 - 6 = 6 | 6 | 36 |
| 13 | 0 | 0 | 13 - 6 = 7 | 0 | 0 |
| 14 | 0 | 0 | 14 - 6 = 8 | 0 | 0 |
| 15 | 0 | 0 | 15 - 6 = 9 | 0 | 0 |
| 16 | 1 | 16 | 16 - 6 = 10 | 10 | 100 |
| Total | $\sum f_i = 40$ | $\sum f_i x_i = 239$ | | $\sum f_i d_i = -1$ | $\sum f_i x_i^2 = 325$ |

$$\therefore \text{Mean } \bar{x} = \frac{\sum f_i x_i}{\sum f_i} = \frac{239}{40} = 5.975 \approx 6$$

and

$$\begin{aligned}
 \sigma &= \sqrt{\frac{\sum f_i d_i^2}{\sum f_i} - \left(\frac{\sum f_i d_i}{\sum f_i}\right)^2} = \sqrt{\frac{325}{40} - \left(\frac{-1}{40}\right)^2} \\
 &= \sqrt{8.125 - 0.000625} = \sqrt{8.124375} = 2.85
 \end{aligned}$$

Q. 19 The weights of coffee in 70 jars is shown in the following table

| Weight (in g) | Frequency |
|---------------|-----------|
| 200-201 | 13 |
| 201-202 | 27 |
| 202-203 | 18 |
| 203-204 | 10 |
| 204-205 | 1 |
| 205-206 | 1 |

Determine variance and standard deviation of the above distribution.

Sol.

| CI | f_i | x_i | $d_i = x_i - \bar{x}$ | $f_i d_i$ | $f_i d_i^2$ |
|---------|-------------------|-------|-----------------------|------------------------|--------------------------|
| 200-201 | 13 | 200.5 | -2 | -26 | 52 |
| 201-202 | 27 | 201.5 | -1 | -27 | 27 |
| 202-203 | 18 | 202.5 | 0 | 0 | 0 |
| 203-204 | 10 | 203.5 | 1 | 10 | 10 |
| 204-205 | 1 | 204.5 | 2 | 2 | 4 |
| 205-206 | 1 | 205.5 | 3 | 3 | 9 |
| | $\Sigma f_i = 70$ | | | $\Sigma f_i d_i = -38$ | $\Sigma f_i d_i^2 = 102$ |

$$\therefore \sigma^2 = \frac{\Sigma f_i d_i^2}{\Sigma f_i} - \left(\frac{\Sigma f_i d_i}{\Sigma f_i} \right)^2 = \frac{102}{70} - \left(\frac{-38}{70} \right)^2$$

Now, $= 1.4571 - 0.2916 = 1.1655$

$$\sigma = \sqrt{1.1655} = 1.08 \text{ g}$$

Q. 20 Determine mean and standard deviation of first n terms of an AP whose first term is a and common difference is d .

Sol.

| x_i | $x_i - a$ | $(x_i - a)^2$ |
|--|-----------|---------------|
| a | 0 | 0 |
| $a + d$ | d | d^2 |
| $a + 2d$ | $2d$ | $4d^2$ |
| | | $9d^2$ |
| | | |
| | | |
| $a + (n-1)d$ | $(n-1)d$ | $(n-1)^2 d^2$ |
| $\Sigma x_i = \frac{n}{2} [2a + (n-1)d]$ | | |

$$\therefore \text{Mean} = \frac{\Sigma x_i}{n} = \frac{1}{n} \left[\frac{n}{2} (2a + (n-1)d) \right]$$

$$= a + \frac{(n-1)}{2} d$$

$$\begin{aligned} \therefore \Sigma(x_i - a) &= d[1 + 2 + 3 + \dots + (n-1)d] \\ &= d \frac{(n-1)n}{2} \end{aligned}$$

$$\begin{aligned} \text{and } \Sigma(x_i - a)^2 &= d^2 [1^2 + 2^2 + 3^2 + \dots + (n-1)^2] \\ &= \frac{d^2(n-1)n(2n-1)}{6} \end{aligned}$$

$$\begin{aligned} \sigma &= \sqrt{\frac{(x_i - a)^2}{n} - \left(\frac{x_i - a}{n}\right)^2} \\ &= \sqrt{\frac{d^2(n-1)n(2n-1)}{6n} - \left[\frac{d(n-1)n}{2n}\right]^2} \\ &= \sqrt{\frac{d^2(n-1)(2n-1)}{6} - \frac{d^2(n-1)^2}{4}} \\ &= d \sqrt{\frac{(n-1)(2n-1)}{6} - \frac{(n-1)^2}{4}} \\ &= d \sqrt{\frac{(n-1)}{2} \left(\frac{2n-1}{3} - \frac{n-1}{2}\right)} \\ &= d \sqrt{\frac{(n-1)}{2} \left[\frac{4n-2-3n+3}{6}\right]} \\ &= d \sqrt{\frac{(n-1)(n+1)}{12}} = d \sqrt{\frac{(n^2-1)}{12}} \end{aligned}$$

Q. 21 Following are the marks obtained, out of 100, by two students Ravi and Hashina in 10 tests

| | | | | | | | | | | |
|----------------|----|----|----|----|----|----|----|----|----|----|
| Ravi | 25 | 50 | 45 | 30 | 70 | 42 | 36 | 48 | 35 | 60 |
| Hashina | 10 | 70 | 50 | 20 | 95 | 55 | 42 | 60 | 48 | 80 |

Who is more intelligent and who is more consistent?

Sol. For Ravi,

| x_i | $d_i = x_i - 45$ | d_i^2 |
|--------------|--------------------|-----------------------|
| 25 | -20 | 400 |
| 50 | 5 | 25 |
| 45 | 0 | 0 |
| 30 | -15 | 225 |
| 70 | 25 | 625 |
| 42 | -3 | 9 |
| 36 | -9 | 81 |
| 48 | 3 | 9 |
| 35 | -10 | 100 |
| 60 | 15 | 225 |
| Total | $\Sigma d_i = -14$ | $\Sigma d_i^2 = 1699$ |

$$\begin{aligned} \sigma &= \sqrt{\frac{\sum d_i^2}{n} - \left(\frac{\sum d_i}{n}\right)^2} \\ &= \sqrt{\frac{1699}{10} - \left(\frac{-14}{10}\right)^2} = \sqrt{169.9 - 0.0196} \\ &= \sqrt{169.88} = 13.03 \end{aligned}$$

Now,

$$\bar{x} = A + \frac{\sum d_i}{\sum f_i} = 45 - \frac{14}{10} = 43.6$$

For Hashina,

| x_i | $d_i = x_i - 55$ | d_i^2 |
|--------------|------------------|---------------------|
| 10 | -45 | 2025 |
| 70 | 25 | 625 |
| 50 | -5 | 25 |
| 20 | -35 | 1225 |
| 95 | 40 | 1600 |
| 55 | 0 | 0 |
| 42 | -13 | 169 |
| 60 | 5 | 25 |
| 48 | -7 | 49 |
| 80 | 25 | 625 |
| Total | $\sum d_i = 0$ | $\sum d_i^2 = 6368$ |

∴ Mean = 55

∴ $\sigma = \sqrt{\frac{6368}{10}} = \sqrt{636.8} = 25.2$

For Ravi, $CV = \frac{\sigma}{\bar{x}} \times 100 = \frac{13.03}{43.6} \times 100 = 29.88$

For Hashina, $CV = \frac{\sigma}{\bar{x}} \times 100 = \frac{25.2}{55} \times 100 = 45.89$

Hence, Hashina is more consistent and intelligent.

Q. 22 Mean and standard deviation of 100 observations were found to be 40 and 10, respectively. If at the time of calculation two observations were wrongly taken as 30 and 70 in place of 3 and 27 respectively, then find the correct standard deviation.

Sol. Given, $n = 100, \bar{x} = 40, \sigma = 10$ and $\bar{x} = 40$

∴ $\frac{\sum x_i}{n} = 40$

⇒ $\frac{\sum x_i}{100} = 40$

⇒ $\sum x_i = 4000$

Now, Corrected $\sum x_i = 4000 - 30 - 70 + 3 + 27$

∴ $= 4030 - 100 = 3930$

Corrected mean = $\frac{3930}{100} = 39.3$

$$\begin{aligned} \text{Now,} \quad \sigma^2 &= \frac{\Sigma x_i^2}{n} - (40)^2 \\ \Rightarrow \quad 100 &= \frac{\Sigma x_i^2}{100} - 1600 \\ \Rightarrow \quad \Sigma x_i^2 &= 170000 \end{aligned}$$

$$\begin{aligned} \text{Now,} \quad \text{Corrected } \Sigma x_i^2 &= 170000 - (30)^2 - (70)^2 + 3^2 + (27)^2 \\ &= 164939 \\ \therefore \quad \text{Corrected } \sigma &= \sqrt{\frac{164939}{100} - (39.3)^2} \\ &= \sqrt{1649.39 - 39.3 \times 39.3} \\ &= \sqrt{1649.39 - 1544.49} \\ &= \sqrt{104.9} = 10.24 \end{aligned}$$

Q. 23 While calculating the mean and variance of 10 readings, a student wrongly used the reading 52 for the correct reading 25. He obtained the mean and variance as 45 and 16, respectively. Find the correct mean and the variance.

$$\begin{aligned} \text{Sol.} \quad \text{Given,} \quad n &= 10, \bar{x} = 45 \text{ and } \sigma^2 = 16 \\ \therefore \quad \bar{x} &= 45 \Rightarrow \frac{\Sigma x_i}{n} = 45 \\ \Rightarrow \quad \frac{\Sigma x_i}{10} &= 45 \Rightarrow \Sigma x_i = 450 \\ \text{Corrected } \Sigma x_i &= 450 - 52 + 25 = 423 \\ \therefore \quad \bar{x} &= \frac{423}{10} = 42.3 \\ \Rightarrow \quad \sigma^2 &= \frac{\Sigma x_i^2}{n} - \left(\frac{\Sigma x_i}{n}\right)^2 \\ \Rightarrow \quad 16 &= \frac{\Sigma x_i^2}{10} - (45)^2 \\ \Rightarrow \quad \Sigma x_i^2 &= 10(2025 + 16) \\ \Rightarrow \quad \Sigma x_i^2 &= 20410 \\ \therefore \quad \text{Corrected } \Sigma x_i^2 &= 20410 - (52)^2 + (25)^2 = 18331 \\ \text{and} \quad \text{corrected } \sigma^2 &= \frac{18331}{10} - (42.3)^2 = 43.81 \end{aligned}$$

Objective Type Questions

Q. 24 The mean deviation of the data 3, 10, 10, 4, 7, 10, 5 from the mean is

- (a) 2 (b) 2.57
(c) 3 (d) 3.75

Sol. (b) Given, observations are 3, 10, 10, 4, 7, 10 and 5.

$$\begin{aligned} \therefore \bar{x} &= \frac{3 + 10 + 10 + 4 + 7 + 10 + 5}{7} \\ &= \frac{49}{7} = 7 \end{aligned}$$

| x_i | $d_i = x_i - \bar{x} $ |
|--------------|-------------------------|
| 3 | 4 |
| 10 | 3 |
| 10 | 3 |
| 4 | 3 |
| 7 | 0 |
| 10 | 3 |
| 5 | 2 |
| Total | $\Sigma d_i = 18$ |

Now,

$$MD = \frac{\Sigma d_i}{N} = \frac{18}{7} = 2.57$$

Q. 25 Mean deviation for n observations x_1, x_2, \dots, x_n from their mean \bar{x} is given by

- (a) $\sum_{i=1}^n (x_i - \bar{x})$ (b) $\frac{1}{n} \sum_{i=1}^n |x_i - \bar{x}|$
(c) $\sum_{i=1}^n (x_i - \bar{x})^2$ (d) $\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$

Sol. (b) $MD = \frac{1}{n} \sum_{i=1}^n |x_i - \bar{x}|$

Q. 26 When tested, the lives (in hours) of 5 bulbs were noted as follows

1357, 1090, 1666, 1494, 1623

The mean deviations (in hours) from their mean is

- (a) 178 (b) 179 (c) 220 (d) 356

Sol. (a) Since, the lives of 5 bulbs are 1357, 1090, 1666, 1494 and 1623.

$$\begin{aligned} \therefore \text{Mean} &= \frac{1357 + 1090 + 1666 + 1494 + 1623}{5} \\ &= \frac{7230}{5} = 1446 \end{aligned}$$

| x_i | $d_i = x_i - \bar{x} $ |
|--------------|-------------------------|
| 1357 | 89 |
| 1090 | 356 |
| 1666 | 220 |
| 1494 | 48 |
| 1623 | 177 |
| Total | $\Sigma d_i = 890$ |

$$MD = \frac{\Sigma d_i}{N} = \frac{890}{5} = 178$$

Q. 27 Following are the marks obtained by 9 students in a mathematics test
50, 69, 20, 33, 53, 39, 40, 65, 59

The mean deviation from the median is

- (a) 9 (b) 10.5
(c) 12.67 (d) 14.76

Sol. (c) Since, marks obtained by 9 students in Mathematics are 50, 69, 20, 33, 53, 39, 40, 65 and 59.

Rewrite the given data in ascending order.

20, 33, 39, 40, 50, 53, 59, 65, 69,

Here,

$$n = 9$$

[odd]

\therefore Median = $\left(\frac{9+1}{2}\right)$ term = 5th term

$$Me = 50$$

| x_i | $d_i = x_i - Me $ |
|---------|--------------------|
| 20 | 30 |
| 33 | 17 |
| 39 | 11 |
| 40 | 10 |
| 50 | 0 |
| 53 | 3 |
| 59 | 9 |
| 65 | 15 |
| 69 | 19 |
| $N = 9$ | $\Sigma d_i = 114$ |

$$\therefore MD = \frac{114}{9} = 12.67$$

Q. 28 The standard deviation of data 6, 5, 9, 13, 12, 8 and 10 is

- (a) $\sqrt{\frac{52}{7}}$ (b) $\frac{52}{7}$
(c) $\sqrt{6}$ (d) 6

$$\begin{aligned} \text{Now,} \quad \sigma &= \sqrt{\frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2} \Rightarrow \sigma^2 = \frac{\sum x_i^2}{n} - (\bar{x})^2 \\ \Rightarrow \quad 25 &= \frac{\sum x_i^2}{100} - (50)^2 \Rightarrow 25 = \frac{\sum x_i^2}{100} - 2500 \\ \Rightarrow \quad 2525 &= \frac{\sum x_i^2}{100} \\ \therefore \quad \sum x_i^2 &= 252500 \end{aligned}$$

Q. 31 If a, b, c, d and e be the observations with mean m and standard deviation s , then find the standard deviation of the observations $a+k, b+k, c+k, d+k$ and $e+k$ is

- (a) s (b) ks (c) $s+k$ (d) $\frac{s}{k}$

Sol. (a) Given observations are a, b, c, d and e .

$$\text{Mean} = m = \frac{a+b+c+d+e}{5}$$

$$\sum x_i = a+b+c+d+e = 5m$$

$$\begin{aligned} \text{Now,} \quad \text{mean} &= \frac{a+k+b+k+c+k+d+k+e+k}{5} \\ &= \frac{(a+b+c+d+e) + 5k}{5} = m+k \end{aligned}$$

$$\begin{aligned} \therefore \quad \text{SD} &= \sqrt{\frac{\sum (x_i+k)^2}{n} - (m+k)^2} \\ &= \sqrt{\frac{\sum (x_i^2 + k^2 + 2kx_i)}{n} - (m^2 + k^2 + 2mk)} \\ &= \sqrt{\frac{\sum x_i^2}{n} - m^2 + \frac{2k\sum x_i}{n} - 2mk} \\ &= \sqrt{\frac{\sum x_i^2}{n} - m^2 + 2km - 2mk} \quad \left[\because \frac{\sum x_i}{n} = m \right] \\ &= \sqrt{\frac{\sum x_i^2}{n} - m^2} \\ &= s \end{aligned}$$

Q. 32 If x_1, x_2, x_3, x_4 and x_5 be the observations with mean m and standard deviation s then, the standard deviation of the observations kx_1, kx_2, kx_3, kx_4 and kx_5 is

- (a) $k+s$ (b) $\frac{s}{k}$ (c) ks (d) s

$$\begin{aligned} \text{Sol. (c)} \quad \text{Here,} \quad m &= \frac{\sum x_i}{5}, s = \sqrt{\frac{\sum x_i^2}{5} - \left(\frac{\sum x_i}{5}\right)^2} \\ \therefore \quad \text{SD} &= \sqrt{\frac{k^2 \sum x_i^2}{5} - \left(\frac{k \sum x_i}{5}\right)^2} \\ &= \sqrt{\frac{k^2 \sum x_i^2}{5} - k^2 \left(\frac{\sum x_i}{5}\right)^2} = \sqrt{\left(\frac{\sum x_i^2}{5}\right) - \left(\frac{\sum x_i}{5}\right)^2} = ks \end{aligned}$$

Q. 33 Let x_1, x_2, \dots, x_n be n observations. Let $w_i = lx_i + k$ for $i = 1, 2, \dots, n$, where l and k are constants. If the mean of x_i 's is 48 and their standard deviation is 12, the mean of w_i 's is 55 and standard deviation of w_i 's is 15, then the value of l and k should be

- (a) $l = 1.25, k = -5$ (b) $l = -1.25, k = 5$
 (c) $l = 2.5, k = -5$ (d) $l = 2.5, k = 5$

Sol. (a) Given, $w_i = lx_i + k, \bar{x}_i = 48, s_{x_i} = 12, \bar{w}_i = 55$ and $s_{w_i} = 15$

Then, $\bar{w}_i = \bar{x}_i + k$ [where, \bar{w}_i is mean w_i 's and \bar{x}_i is mean of x_i 's]
 $\Rightarrow 55 = 48 + k$... (i)

Now, SD of $w_i =$ SD of x_i
 $\Rightarrow 15 = 12$
 $\Rightarrow l = \frac{15}{12} = 1.25$... (ii)

From Eqs. (i) and (ii), $k = 55 - 1.25 \times 48 = -5$

Q. 34 The standard deviations for first natural numbers is

- (a) 5.5 (b) 3.87 (c) 2.97 (d) 2.87

Sol. (d) We know that, SD of first n natural number = $\sqrt{\frac{n^2 - 1}{12}}$

\therefore SD of first 10 natural numbers = $\sqrt{\frac{(10)^2 - 1}{12}} = \sqrt{\frac{100 - 1}{12}} = \sqrt{\frac{99}{12}} = \sqrt{8.25} = 2.87$

Q. 35 Consider the numbers 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10. If 1 is added to each number the variance of the numbers, so obtained is

- (a) 6.5 (b) 2.87 (c) 3.87 (d) 8.25

Sol. (d) Given numbers are 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10.

If 1 is added to each number, then observations will be 2, 3, 4, 5, 6, 7, 8, 9, 10 and 11.

$\therefore \Sigma x_i = 2 + 3 + 4 + \dots + 11 = \frac{10}{2} [2 \times 2 + 9 \times 1] = 5[4 + 9] = 65$

and $\Sigma x_i^2 = 2^2 + 3^2 + 4^2 + 5^2 + \dots + 11^2 = (1^2 + 2^2 + 3^2 + \dots + 11^2) - (1^2) = \frac{11 \times 12 \times 23}{6} - 1 = \frac{11 \times 12 \times 23 - 6}{6} = 505$

$$\begin{aligned} \therefore s^2 &= \frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2 = \frac{505}{10} - \left(\frac{65}{10}\right)^2 \\ &= 50.5 - (6.5)^2 \\ &= 50.5 - 42.25 \\ &= 8.25 \end{aligned}$$

Q. 36 Consider the first 10 positive integers. If we multiply each number by -1 and, then add 1 to each number, the variance of the numbers, so obtained is

- (a) 8.25 (b) 6.5 (c) 3.87 (d) 2.87

Sol. (a) Since, the first 10 positive integers are 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10.

On multiplying each number by -1 , we get

$$-1, -2, -3, -4, -5, -6, -7, -8, -9, -10$$

On adding 1 in each number, we get

$$0, -1, -2, -3, -4, -5, -6, -7, -8, -9$$

$$\begin{aligned} \therefore \sum x_i &= 0 - 1 - 2 - 3 - 4 - 5 - 6 - 7 - 8 - 9 \\ &= -\frac{9 \times 10}{2} \end{aligned}$$

$$= -45$$

$$\begin{aligned} \text{and } \sum x_i^2 &= 0^2 + (-1)^2 + (-2)^2 + \dots + (-9)^2 \\ &= \frac{9 \times 10 \times 19}{6} \end{aligned}$$

$$= 285$$

$$\begin{aligned} \therefore \text{SD} &= \sqrt{\frac{285}{10} - \left(\frac{-45}{10}\right)^2} = \sqrt{\frac{285}{10} - \frac{2025}{100}} \\ &= \sqrt{\frac{2850 - 2025}{100}} = \sqrt{8.25} \end{aligned}$$

$$\text{Now, variance} = (\text{SD})^2 = (\sqrt{8.25})^2 = 8.25$$

Q. 37 The following information relates to a sample of size 60, $\sum x^2 = 18000$, and $\sum x = 960$. Then, the variance is

- (a) 6.63 (b) 16 (c) 22 (d) 44

$$\begin{aligned} \text{Sol. (d)} \quad \text{Variance} &= \frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2 \\ &= \frac{18000}{60} - \left(\frac{960}{60}\right)^2 = 300 - 256 = 44 \end{aligned}$$

Q. 38 If the coefficient of variation of two distributions are 50, 60 and their arithmetic means are 30 and 25 respectively, then the difference of their standard deviation is

- (a) 0 (b) 1
(c) 1.5 (d) 2.5

Sol. (a) Here $CV_1 = 50$, $CV_2 = 60$, $\bar{x}_1 = 30$ and $\bar{x}_2 = 25$

$$\begin{aligned} \therefore CV_1 &= \frac{\sigma_1}{\bar{x}_1} \times 100 \Rightarrow 50 = \frac{\sigma_1}{30} \times 100 \\ \therefore \sigma_1 &= \frac{30 \times 50}{100} = 15 \text{ and } CV_2 = \frac{\sigma_2}{\bar{x}_2} \times 100 \\ \Rightarrow 60 &= \frac{\sigma_2}{25} \times 100 \\ \therefore \sigma_2 &= \frac{60 \times 25}{100} = 15 \\ \text{Now, } \sigma_1 - \sigma_2 &= 15 - 15 = 0 \end{aligned}$$

Q. 39 The standard deviation of some temperature data in °C is 5. If the data were converted into °F, then the variance would be

- (a) 81 (b) 57 (c) 36 (d) 25

Sol. (a) Given, $\sigma_C = 5 \Rightarrow \frac{5}{9}(F - 32) = C$

$$F = \frac{9C}{5} + 32$$

$$\sigma_F = \frac{9}{5} \sigma_C = \frac{9}{5} \times 5 = 9$$

Here, $\sigma_F^2 = (9)^2 = 81$

Fillers

Q. 40 Coefficient of variation = $\frac{\dots}{\text{Mean}} \times 100$

Sol. $CV = \frac{SD}{\text{Mean}} \times 100$

Q. 41 If \bar{x} is the mean of n values of x , then $\sum_{i=1}^n (x_i - \bar{x})$ is always equal to

..... . If a has any value other than \bar{x} , then $\sum_{i=1}^n (x_i - \bar{x})^2$ is than $\sum (x_i - a)^2$

Sol. If \bar{x} is the mean of n values of x , then $\sum_{i=1}^n (x_i - \bar{x}) = 0$ and if a has any value other than \bar{x} , then

$$\sum_{i=1}^n (x_i - \bar{x})^2 \text{ is less than } \sum (x_i - a)^2.$$

Q. 42 If the variance of a data is 121, then the standard deviation of the data is

Sol. If the variance of a data is 121.

Then,

$$\begin{aligned} \text{SD} &= \sqrt{\text{Variance}} \\ &= \sqrt{121} = 11 \end{aligned}$$

Q. 43 The standard deviation of a data is of any change in origin but is of change of scale.

Sol. The standard deviation of a data is independent of any change in origin but is dependent of change of scale.

Q. 44 The sum of squares of the deviations of the values of the variable is when taken about their arithmetic mean.

Sol. The sum of the squares of the deviations of the values of the variable is minimum when taken about their arithmetic mean.

Q. 45 The mean deviation of the data is when measured from the median.

Sol. The mean deviation of the data is least when measured from the median.

Q. 46 The standard deviation is to the mean deviation taken from the arithmetic mean.

Sol. The SD is greater than or equal to the mean deviation taken from the arithmetic mean.