CIRCLES

EXERCISE 10.1

Q.1.	Fill in the blanks:
	(i) The centre of a circle lies in of the circle. (exterior) interior)
	(ii) A point, whose distance from the centre of a circle is greater than its radius lies in of the circle. (exterior/interior)
	(iii) The longest chord of a circle is a of the circle.
	(iv) An arc is a when its ends are the ends of a diameter.
	(v) Segment of a circle is the region between an arc and o _i the circle.
	(vi) A circle divides the plane, on which it lies in parts.
Sol.	(i) interior (ii) exterior (iii) diameter (iv) semicircle (v) the chord (vi
	three
Q.2.	Write True or False: Give reasons for your answers.
	(i) Line segment joining the centre to any point on the circle is a radius of the circle.
	(ii) A circle has only finite number of equal chords.
	(iii) If a circle is divided into three equal arcs, each is a major arc.
	(iv) A chord of a circle, which is twice as long as its radius, is a diameter of the circle.
	(v) Sector is the region between the chord and its corresponding arc.
	(vi) A circle is a plane figure.
Sol.	(i) True (ii) False (iii) False (iv) True (v) False (vi) True


CIRCLES

EXERCISE 10.2


- **Q.1.** Recall that two circles are congruent if they have the same radii. Prove that equal chords of congruent circles subtend equal angles at their centres.
- **Sol. Given :** Two congruent circles with centres O and O'. AB and CD are equal chords of the circles with centres O and O' respectively.

To Prove : ∠AOB = ∠COD


Proof: In triangles AOB and COD,



 $\Rightarrow \angle AOB \cong \angle CO'D$ **Proved.** [CPCT]



Ans. Given: Two congruent circles with centres O and O'. AB and CD are chords of circles with centre O and O' respectively such that ∠AOB = ∠CO'D





Proof: In triangles AOB and CO'D,

$$AO = CO'$$
 $BO = DO'$
 $AOB = \angle CO'D$ [Given]

 $AOB = \angle CO'D$ [SAS axiom]

 \Rightarrow AB = CD **Proved.** [CPCT]





CIRCLES


EXERCISE 10.3

Q.1. Draw different pairs of circles. How many points does each pair have in common? What is the maximum number of common points?


Ans.



(i) 0 point



(ii) 1 point




(iii) 2 points

Maximum number of common points = 2 Ans.

- **Q.2.** Suppose you are given a circle. Give a construction to find its centre.
- Ans. Steps of Construction:
 - 1. Take arc PQ of the given circle.
 - 2. Take a point R on the arc PQ and draw chords PR and RQ.
 - 3. Draw perpendicular bisectors of PR and RQ. These perpendicular bisectors intersect at point O.

Hence, point O is the centre of the given circle.



Ans. Given: AB is the common chord of two intersecting circles (O, r) and (O', r'). **To Prove:** Centres of both circles lie on the perpendicular bisector of chord AB, i.e., AB is bisected at right angle by OO'.

Construction: Join AO, BO, AO' and BO'.

Proof: In $\triangle AOO'$ and $\triangle BOO'$

$$AO = OB$$
 (Radii of the circle (O, r))
 $AO' = BO'$ (Radii of the circle (O', r'))

$$OO' = OO'$$
 (Common)

$$\triangle \Delta AOO' \cong \Delta BOO'$$
 (SSS congruency)


$$\Rightarrow \angle AOO' = \angle BOO' (CPCT)$$

Now in $\triangle AOC$ and $\triangle BOC$

$$\angle AOC = \angle BOC \quad (\angle AOO' = \angle BOO')$$

AO = BO (Radii of the circle
$$(O, r)$$
)

$$OC = OC$$
 (Common)



```
∴ \triangle AOC \cong \triangle BOC (SAS congruency)

⇒ AC = BC and \angle ACO = \angle BCO ...(i) (CPCT)

⇒ \angle ACO + \angle BCO = 180^{\circ} ...(ii) (Linear pair)

⇒ \angle ACO = \angle BCO = 90^{\circ} (From (i) and (ii))

Hence, OO' lie on the perpendicular bisector of AB
```

MMM. Greathing opposition

CIRCLES

EXERCISE 10.4

- **Q.1.** Two circles of radii 5 cm and 3 cm intersect at two points and the distance between their centres is 4 cm. Find the length of the common chord.
- **Sol.** In $\triangle AOO'$,

$$AO^{2} = 5^{2} = 25$$

 $AO'^{2} = 3^{2} = 9$
 $OO'^{2} = 4^{2} = 16$
 $AO'^{2} + OO'^{2} = 9 + 16 = 25 = AO^{2}$
 $\Rightarrow \angle AO'O$
 $= 90^{\circ}$

[By converse of pythagoras theorem]

Similarly, $\angle BO'O = 90^\circ$.

$$\Rightarrow \angle AO'B = 90^{\circ} + 90^{\circ} = 180^{\circ}$$

$$\Rightarrow$$
 AO'B is a straight line. whose mid-point is O.

$$\Rightarrow$$
 AB = $(3 + 3)$ cm = 6 cm **Ans.**

- **Q.2.** If two equal chords of a circle intersect within the circle, prove that the segments of one chord are equal to corresponding segments of the other chord.
- **Sol. Given:** AB and CD are two equal chords of a circle which meet at E. **To prove:** AE = CE and BE = DE

Construction : Draw OM \perp AB and ON \perp CD and join OE. **Proof :** In Δ OME and Δ ONE

$$OE = OE$$
 [Common]

 \angle OME = \angle ONE [Each equal to 90°]

$$\therefore \quad \Delta OME \cong \Delta ONE \qquad [RHS axiom]$$

$$\Rightarrow \qquad \text{EM} = \text{EN} \qquad \dots \text{(i)} \qquad \text{[CPCT]}$$

Now
$$AB = CD$$
 [Given]

$$\Rightarrow \quad \frac{1}{2} AB = \frac{1}{2} CD$$

$$\Rightarrow$$
 AM = CN ...(ii) [Perpendicular from

centre bisects the chord]

$$EM + AM = EN + CN$$


$$\Rightarrow \qquad AE = CE \qquad ..(iii)$$

Now,
$$AB = CD$$
 ...(iv)
 $\Rightarrow AB - AE = CD - AE$ [From (iii)]

$$\Rightarrow$$
 BE = CD - CE **Proved.**

- **Q.3.** If two equal chords of a circle intersect within the circle, prove that the line joining the point of intersection to the centre makes equal angles with the chords.
- **Sol.** Given: AB and CD are two equal chords of a circle which meet at E within the circle and a line PQ joining the point of intersection to the centre.

To Prove : ∠AEQ = ∠DEQ



Construction : Draw $OL \perp AB$ and $OM \perp CD$.

Proof: In \triangle OLE and \triangle OME, we have

OL = OM [Equal chords are equidistant]

$$OE = OE$$

[Common]

 $[Each = 90^{\circ}]$

$$\therefore \Delta OLE \cong \Delta OME$$

[RHS congruence]

[CPCT]



Q.4. If a line intersects two concentric circles (circles with the same centre) with centre O at A, B, C and D, prove that AB = CD (see Fig.)

Sol. Given: A line AD intersects two concentric circles at A, B, C and D, where O is the centre of these circles.

To prove : AB = CD

Construction : Draw OM \perp AD.

Proof: AD is the chord of larger circle.

:. AM = DM ..(i) [OM bisects the chord]


BC is the chord of smaller circle


 \therefore BM = CM ...(ii) [OM bisects the chord]

Subtracting (ii) from (i), we get

$$AM - BM = DM - CM$$

$$\Rightarrow$$
 AB = CD **Proved.**





Draw OL
$$\perp$$
 RS,

$$OL^2 = OR^2 - RL^2$$

$$OL^2 = 5^2 - 3^2$$
 [RL = 3 m, because OL \perp RS]
= 25 - 9 = 16

$$OL = \sqrt{16} = 4$$

Now, area of triangle ORS = $\frac{1}{2} \times KR \times 05$

$$= \frac{1}{2} \times KR \times 05$$

Also, area of $\triangle ORS = \frac{1}{2} \times RS \times OL = \frac{1}{2} \times 6 \times 4 = 12 \text{ m}^2$


$$\Rightarrow \frac{1}{2} \times KR \times 5 = 12$$

$$\Rightarrow$$
 KR = $\frac{12 \times 2}{5} = \frac{24}{5} = 4.8 \text{ m}$

$$\Rightarrow$$
 RM = 2KR

$$\Rightarrow$$
 RM = 2 × 4.8 = 9.6 m

Hence, distance between Reshma and Mandip is 9.6 m Ans.



- **Q.6.** A circular park of radius 20 m is situated in a colony. Three boys Ankur, Syed and David are siting at equal distance on its boundary each having a toy telephone in his hands to talk each other. Find the length of the string of each phone.
- **Sol.** Let Ankur, Syed and David be represented by A, S and D respectively.

Let PD = SP = SQ = QA = AR = RD =
$$x$$
 In \triangle OPD,

$$OP^2 = 400 - x^2$$

$$\Rightarrow$$
 OP = $\sqrt{400-x^2}$

⇒ AP =
$$2\sqrt{400 - x^2} + \sqrt{400 - x^2}$$

[: centroid divides the median in the ratio 2 : 1]

$$= 3\sqrt{400 - x^2}$$

Now, in $\triangle APD$,

$$PD^2 = AD^2 - DP^2$$

$$\Rightarrow x^2 = (2x)^2 - (3\sqrt{400 - x^2})^2$$

$$\Rightarrow$$
 $x^2 = 4x^2 - 9(400 - x^2)$

$$\Rightarrow$$
 $x^2 = 4x^2 - 3600 + 9x^2$

$$\Rightarrow$$
 12 x^2 = 3600

$$\Rightarrow \qquad x^2 = \frac{3600}{12} = 300$$

$$\Rightarrow$$
 $x = 10\sqrt{3}$

Now, SD =
$$2x = 2 \times 10\sqrt{3} = 20\sqrt{3}$$

∴ ASD is an equilateral triangle.


$$\Rightarrow$$
 SD = AS = AD = $20\sqrt{3}$

Hence, length of the string of each phone is $20 \sqrt{3}$ m Ans.

CIRCLES

EXERCISE 10.5

Q.1. In the figure, A, B and C are three points on a circle with centre O such that $\angle BOC = 30^{\circ}$ and $\angle AOB = 60^{\circ}$. If D is a point on the circle other than the arc ABC, find $\angle ADC$.



Sol. We have,
$$\angle BOC = 30^{\circ}$$
 and $\angle AOB = 60^{\circ}$
 $\angle AOC = \angle AOB + \angle BOC = 60^{\circ} + 30^{\circ} = 90^{\circ}$

$$\Rightarrow \angle ADC = \frac{1}{2} \angle AOC = \frac{1}{2} \times 90^{\circ} \Rightarrow \angle ADC = 45^{\circ} \text{ Ans.}$$

- **Q.2.** A chord of a circle is equal to the radius of the circle. Find the angle subtended by the chord at a point on the minor arc and also at a point on the major arc.
- **Sol.** We have, OA = OB = AB

Therefore, $\triangle OAB$ is a equilateral triangle.

$$\Rightarrow$$
 $\angle AOB = 60^{\circ}$


We know that angle subtended by an arc at the centre of a circle is double the angle subtended by the same arc on the remaining part of the circle.

$$\Rightarrow$$
 $\angle ACB = \frac{1}{2} \angle AOB = \frac{1}{2} \times 60^{\circ}$

$$\Rightarrow$$
 $\angle ACB = 30^{\circ}$

Also,
$$\angle ADB = \frac{1}{2} \text{ reflex } \angle AOB$$

$$= \frac{1}{2}(360^{\circ} - 60^{\circ}) = \frac{1}{2} \times 300^{\circ} = 150^{\circ}$$




Hence, angle subtended by the chord at a point on the minor arc is 150° and at a point on the major arc is 30° **Ans.**

Q.3. In the figure, $\angle PQR = 100^{\circ}$, where P, Q and R are points on a circle with centre O. Find $\angle OPR$.

$$=2\times100^{\circ}=200^{\circ}$$

Now, angle POR =
$$360^{\circ} - 200 = 160^{\circ}$$
 Also,



$$\angle OPR = \angle ORP$$
 [Opposite angles of isosceles triangle] In $\triangle OPR$, $\angle POR = 160^{\circ}$

$$\therefore$$
 $\angle OPR = \angle ORP = 10^{\circ}$

[Angle sum property of a triangle]. Ans.

Q.4. In the figure,
$$\angle ABC = 69^{\circ}$$
, $\angle ACB = 31^{\circ}$, find $\angle BDC$.

Sol. In
$$\triangle$$
ABC, we have

$$\angle ABC + \angle ACB + \angle BAC = 180^{\circ}$$


[Angle sum property of a triangle]

$$\Rightarrow$$
 69° + 31° + \angle BAC = 180°

$$\Rightarrow$$
 $\angle BAC = 180^{\circ} - 100^{\circ} = 80^{\circ}$

Also, $\angle BAC = \angle BDC$ [Angles in the same segment]

$$\Rightarrow$$
 $\angle BDC = 80^{\circ}$ Ans.



Q.5. In the figrue, A, B, C and D are four points on a circle. AC and BD intersect at a point E such that ∠BEC = 130° and ∠ECD = 20°. Find ∠BAC.

Sol.
$$\angle BEC + \angle DEC = 180^{\circ}$$
 [Linear pair]

$$\Rightarrow$$
 130° + \angle DEC = 180°

$$\Rightarrow$$
 $\angle DEC = 180^{\circ} - 130^{\circ} = 50^{\circ}$

Now, in $\triangle DEC$,

$$\Rightarrow$$
 \angle DEC + \angle DCE + \angle CDE = 180°


[Angle sum property of a triangle]

$$\Rightarrow 50^{\circ} + 20^{\circ} + \angle CDE = 180^{\circ}$$

$$\Rightarrow$$
 $\angle CDE = 180^{\circ} - 70^{\circ} = 110^{\circ}$

Also, $\angle CDE = \angle BAC$ [Angles in same segment]

$$\Rightarrow$$
 $\angle BAC = 110^{\circ} \text{ Ans.}$



- **Q.6.** ABCD is a cyclic quadrilateral whose diagonals intersect at a point E. If $\angle DBC = 70^{\circ}$, $\angle BAC = 30^{\circ}$, find $\angle BCD$. Further, if AB = BC, find $\angle ECD$.
- **Sol.** $\angle CAD = \angle DBC = 70^{\circ}$ [Angles in the same segment] Therefore, $\angle DAB = \angle CAD + \angle BAC$

$$= 70^{\circ} + 30^{\circ} = 100^{\circ}$$

But, $\angle DAB + \angle BCD = 180^{\circ}$

[Opposite angles of a cyclic quadrilateral]

So,
$$\angle BCD = 180^{\circ} - 100^{\circ} = 80^{\circ}$$

Now, we have AB = BC

Therefore, $\angle BCA = 30^{\circ}$ [Opposite angles of an isosceles triangle] Again, $\angle DAB + \angle BCD = 180^{\circ}$

[Opposite angles of a cyclic quadrilateral]

$$\Rightarrow 100^{\circ} + \angle BCA + \angle ECD = 180^{\circ} \ [\because \angle BCD = \angle BCA + \angle ECD]$$

$$\Rightarrow 100^{\circ} + 30^{\circ} + \angle ECD = 180^{\circ}$$

$$\Rightarrow$$
 130° + \angle ECD = 180°

$$\Rightarrow \angle ECD = 180^{\circ} - 130^{\circ} = 50^{\circ}$$

Hence, $\angle BCD = 80^{\circ}$ and $\angle ECD = 50^{\circ}$ Ans.

- **Q.7.** If diagonals of a cyclic quadrilateral are diameters of the circle through the vertices of the quadrilateral, prove that it is a rectangle.
- **Sol. Given :** ABCD is a cyclic quadrilateral, whose diagonals AC and BD are diameter of the circle passing through A, B, C and D.



Proof : In $\triangle AOD$ and $\triangle COB$

$$OD = OB$$
 [Radii of a circle]
 $\angle AOD = \angle COB$ [Vertically opposite angles]

$$\therefore \quad \Delta AOD \cong \Delta COB \quad [SAS axiom]$$

$$\therefore \quad \angle OAD = \angle OCB \quad [CPCT]$$

But these are alternate interior angles made by the transversal AC, intersecting AD and BC.

$$\therefore$$
 AD \parallel BC

Similarly, AB || CD.

Hence, quadrilateral ABCD is a parallelogram.

Also,
$$\angle ABC = \angle ADC$$
 ..(i) [Opposite angles of a ||gm are equal]

And,
$$\angle ABC + \angle ADC = 180^{\circ}$$
 ...(ii)

[Sum of opposite angles of a cyclic quadrilateral is 180°]

$$\Rightarrow \angle ABC = \angle ADC = 90^{\circ}$$
 [From (i) and (ii)]

$$\therefore$$
 ABCD is a rectangle. $\;$ [A $\parallel\!\text{gm}$ one of whose angles is

90° is a rectangle] **Proved.**


- Q.8. If the non-parallel sides of a trapezium are equal, prove that it is cyclic.
- **Sol.** Given: A trapezium ABCD in which AB || CD and AD = BC.

To Prove: ABCD is a cyclic trapezium.

Construction : Draw DE \perp AB and CF \perp AB.

Proof: In $\triangle DEA$ and $\triangle CFB$, we have

$$\angle DEA = \angle CFB = 90^{\circ} [DE \perp AB \text{ and } CF \perp AB]$$



DE = CF

[Distance between parallel lines remains constant]

∴
$$\Delta DEA \cong \Delta CFB$$
 [RHS axiom]
⇒ $\angle A = \angle B$...(i) [CPCT]

and,
$$\angle ADE = \angle BCF$$
 ..(ii) [CPCT]

Since,
$$\angle ADE = \angle BCF$$
 [From (ii)]

$$\Rightarrow$$
 $\angle ADE + 90^{\circ} = \angle BCF + 90^{\circ}$

$$\Rightarrow \angle ADE + \angle CDE = \angle BCF + \angle DCF$$

$$\Rightarrow$$
 $\angle D = \angle C$...(iii)

$$[\angle ADE + \angle CDE = \angle D, \angle BCF + \angle DCF = \angle C]$$

$$\therefore \angle A = \angle B \text{ and } \angle C = \angle D$$
 [From (i) and (iii)] (iv)

$$\angle A + \angle B + \angle C + \angle D = 360^{\circ}$$
 [Sum of the angles of a quadrilateral is 360°]

$$\Rightarrow 2(\angle B + \angle D) = 360^{\circ}$$
 [Using (iv)]

$$\Rightarrow \angle B + \angle D = 180^{\circ}$$

⇒ Sum of a pair of opposite angles of quadrilateral ABCD is 180°.

$$\Rightarrow$$
 ABCD is a cyclic trapezium **Proved.**

- **Q.9.** Two circles intersect at two points B and C. Through B, two line segments ABD and PBQ are drawn to intersect the circles at A, D and P, Q respectively (see Fig.). Prove that $\angle ACP = \angle QCD$.
- **Sol. Given:** Two circles intersect at two points B and C. Through B, two line segments ABD and PBQ are drawn to intersect the circles at A, D and P, Q respectively.

To Prove :
$$\angle$$
ACP = \angle QCD.

Proof:
$$\angle ACP = \angle ABP$$
 ...(i)

$$\angle QCD = \angle QBD$$
 ..(ii)

[Angles in the same segment]

$$\angle ACP = \angle QCD$$
 Proved.

- **Q.10.** If circles are drawn taking two sides of a triangle as diameters, prove that the point of intersection of these circles lie on the third side.
 - **Sol. Given :** Sides AB and AC of a triangle ABC are diameters of two circles which intersect at D.





Also,
$$\angle ADC = 90^{\circ}$$
 ..(ii)

Adding (i) and (ii), we get


$$\angle ADB + \angle ADC = 90^{\circ} + 90^{\circ}$$

$$\Rightarrow$$
 $\angle ADB + \angle ADC = 180^{\circ}$

- \Rightarrow BDC is a straight line.
- : D lies on BC

Hence, point of intersection of circles lie on the third side BC. Proved.


- **Q.11.** ABC and ADC are two right triangles with common hypotenuse AC. Prove that $\angle CAD = \angle CBD$.
 - **Sol. Given :** ABC and ADC are two right triangles with common hypotenuse AC. **To Prove :** ∠CAD = ∠CBD



Proof: Let O be the mid-point of AC.


Then OA = OB = OC = OD

Mid point of the hypotenuse of a right triangle is equidistant from its vertices with O as centre and radius equal to OA, draw a circle to pass through A, B, C and D.



We know that angles in the same segment of a circle are equal.

Since, $\angle CAD$ and $\angle CBD$ are angles of the same segment. Therefore, $\angle CAD = \angle CBD$. **Proved.**




Sol. Given : ABCD is a cyclic parallelogram.

To prove : ABCD is a rectangle.

Proof: $\angle ABC = \angle ADC$...(i)

[Opposite angles of a ||gm are equal]

But, $\angle ABC + \angle ADC = 180^{\circ}$... (ii)




$$\Rightarrow \angle ABC = \angle ADC = 90^{\circ}$$
 [From (i) and (ii)]

∴ ABCD is a rectangle

[A ||gm one of whose angles is 90° is a rectangle]

Hence, a cyclic parallelogram is a rectangle. **Proved.**



CIRCLES

EXERCISE 10.6 (Optional)

- Q.1. Prove that the line of centres of two intersecting circles subtends equal angles at the two points of intersection.
- **Sol. Given:** Two intersecting circles, in which OO' is the line of centres and A and B are two points of intersection.

To prove : ∠OAO′ = ∠OBO′

Construction : Join AO, BO, AO' and BO'. **Proof :** In ΔAOO' and ΔBOO', we have

AO = BO [Radii of the same circle] AO' = BO' [Radii of the same circle]

OO' = OO' [Common] $\triangle AOO' \cong \triangle BOO'$ [SSS axiom] $\angle OAO' = \angle OBO'$ [CPCT]

Hence, the line of centres of two intersecting circles subtends equal angles at the two points of intersection. **Proved.**


- **Q.2.** Two chords AB and CD of lengths 5 cm and 11 cm respectively of a circle are parallel to each other and are on opposite sides of its centre. If the distance between AB and CD is 6 cm, find the radius of the circle.
- **Sol.** Let O be the centre of the circle and let its radius be r cm.

Draw OM \perp AB and OL \perp CD.

Then, AM =
$$-\frac{1}{2}$$
AB = $\frac{5}{2}$ cm

and,
$$CL = \frac{1}{2}CD = \frac{11}{2}$$
 cm

Since, AB \parallel CD, it follows that the points O, L, M are



collinear and therefore, LM = 6 cm.

Let OL = x cm. Then OM = (6 - x) cm

Join OA and OC. Then OA = OC = r cm.

Now, from right-angled Δ OMA and Δ OLC, we have

 $OA^2 = OM^2 + AM^2$ and $OC^2 = OL^2 + CL^2$ [By Pythagoras Theorem]

$$\Rightarrow r^2 = (6 - x)^2 + \left(\frac{5}{2}\right)^2$$
 ...(i) and $r^2 = x^2 + \left(\frac{11}{2}\right)^2$... (ii)

$$\Rightarrow (6 - x)^{2} + \left(\frac{5}{2}\right)^{2} = x^{2} + \left(\frac{11}{2}\right)^{2} \text{ [From (i) and (ii)]}$$

$$\Rightarrow 36 + x^{2} - 12x + \frac{25}{4} = x^{2} + \frac{121}{4}$$

$$\Rightarrow -12x = \frac{121}{4} - \frac{25}{4} - 36$$

$$\Rightarrow -12x = \frac{96}{4} - 36$$

$$\Rightarrow -12x = 24 - 36$$

$$\Rightarrow -12x = -12$$

$$\Rightarrow x = 1$$
So that the first second of the content of the

Substituting x = 1 in (i), we get

$$r^{2} = (6 - x)^{2} + \left(\frac{5}{2}\right)^{2}$$

$$\Rightarrow r^{2} = (6 - 1)^{2} + \left(\frac{5}{2}\right)^{2}$$

$$\Rightarrow r^{2} = (5)^{2} + \left(\frac{5}{2}\right)^{2} = 25 + \frac{25}{4}$$

$$\Rightarrow r^{2} = \frac{125}{4}$$


$$\Rightarrow r = \frac{5\sqrt{5}}{2}$$

Hence, radius $r = \frac{5\sqrt{5}}{2}$ cm. Ans.

- **Q.3.** The lengths of two parallel chords of a circle are 6 cm and 8 cm. If the smaller chord is at distance 4 cm from the centre, what is the distance of the other chord from the centre?
- **Sol.** Let PQ and RS be two parallel chords of a circle with centre O. We have, PQ = 8 cm and RS = 6 cm.

Draw perpendicular bisector OL of RS which meets PQ in M. Since, PQ || RS, therefore, OM is also perpendicular bisector of PQ.

Also, OL = 4 cm and RL =
$$\frac{1}{2}$$
 RS \Rightarrow RL = 3 cm and PM = $\frac{1}{2}$ PQ \Rightarrow PM = 4 cm In \triangle ORL, we have OR² = RL² + OL² [Pythagoras theorem]



$$\Rightarrow$$
 OR² = 3² + 4² = 9 + 16

$$\Rightarrow$$
 OR² = 25 \Rightarrow OR = $\sqrt{25}$

$$\Rightarrow$$
 OR = 5 cm

$$\therefore$$
 OR = OP

[Radii of the circle]

$$\Rightarrow$$
 OP = 5 cm

Now, in $\triangle OPM$

$$OM^2 = OP^2 - PM^2$$
 [Pythagoras theorem]

$$\Rightarrow$$
 OM² = 5² - 4² = 25 - 16 = 9

$$OM = \sqrt{9} = 3 \text{ cm}$$

Hence, the distance of the other chord from the centre is 3 cm. Ans.

- **Q.4.** Let the vertex of an angle ABC be located outside a circle and let the sides of the angle intersect equal chords AD and CE with the circle. Prove that ∠ABC is equal to half the difference of the angles subtended by the chords AC and DE at the centre.
- **Sol. Given :** Two equal chords AD and CE of a circle with centre O. When meet at B when produced.

To Prove :
$$\angle ABC = \frac{1}{2}(\angle AOC - \angle DOE)$$

Proof: Let
$$\angle AOC = x$$
, $\angle DOE = y$, $\angle AOD = z$

$$\angle EOC = z$$
 [Equal chords subtends equal angles at the centre]

[Angle at a point]

$$\therefore x + y + 2z = 36^{\circ}$$

$$OA = OD \implies \angle OAD = \angle ODA$$

$$\angle OAD + \angle ODA + z = 180^{\circ}$$

$$\Rightarrow 2\angle OAD = 180^{\circ} - z \qquad [\because \angle OAD = \angle OBA]$$

$$\Rightarrow \angle OAD = 90^{\circ} - \frac{z}{2}$$
 ... (ii)

Similarly
$$\angle OCE = 90^{\circ} - \frac{z}{2}$$
 ... (iii)

$$\Rightarrow \angle ODB = \angle OAD + \angle ODA$$
 [Exterior angle property]

$$\Rightarrow \angle OEB = 90^{\circ} - \frac{z}{2} + z$$
 [From (ii)]

$$\Rightarrow \angle \text{ODB} = 90^{\circ} + \frac{z}{2}$$
 ... (iv)

Also,
$$\angle OEB = \angle OCE + \angle COE$$
 [Exterior angle property]

$$\Rightarrow \angle OEB = 90^{\circ} - \frac{z}{2} + z$$
 [From (iii)]

$$\Rightarrow \angle OEB = 90^{\circ} + \frac{z}{2}$$
 ... (v)

Also,
$$\angle OED = \angle ODE = 90^{\circ} - \frac{y}{2}$$
 ... (vi)

O from (iv), (v) and (vi), we have

$$\angle BDE = \angle BED = 90^{\circ} + \frac{z}{2} - \left(90^{\circ} - \frac{y}{2}\right)$$

$$\Rightarrow \angle BDE = \angle BED = \frac{y+z}{2}$$

$$\Rightarrow \angle BDE = \angle BED = y + z$$
 ... (vii)

$$\therefore$$
 \angle BDE = $180^{\circ} - (y + z)$

$$\Rightarrow \angle ABC = 180^{\circ} - (y + z)$$
 ... (viii)

Now,
$$\frac{y-z}{2} = \frac{360^{\circ} - y - 2z - y}{2} = 180^{\circ} - (y+z)$$
 ... (ix)

From (viii) and (ix), we have


$$\angle ABC = \frac{x-y}{2}$$
 Proved.

- **Q.5.** Prove that the circle drawn with any side of a rhombus as diameter, passes through the point of intersection of its diagonals.
- **Sol. Given:** A rhombus ABCD whose diagonals intersect each other at O. **To prove:** A circle with AB as diameter passes through O.

Proof :
$$\angle AOB = 90^{\circ}$$

[Diagonals of a rhombus bisect each other at 90°]

- \Rightarrow \triangle AOB is a right triangle right angled at O.
- \Rightarrow AB is the hypotenuse of A B right \triangle AOB.
- ⇒ If we draw a circle with AB as diameter, then it will pass through O. because angle is a semicircle is 90° and $\angle AOB = 90^{\circ}$ **Proved.**



(ii)

- **Q.6.** ABCD is a parallelogram. The circle through A, B and C intersect CD (produced if necessary) at E. Prove that AE = AD.
- **Sol. Given :** ABCD is a parallelogram.

Construction: Draw a circle which passes through ABC and intersect CD (or CD produced) at E.

$$\angle AED + \angle ABC = 180^{\circ}$$

But
$$\angle ACD = \angle ADC = \angle ABC + \angle ADE$$

$$\Rightarrow$$
 $\angle ABC + \angle ADE = 180^{\circ}$ [From (ii)] ... (iii)


From (i) and (iii)

$$\angle AED + \angle ABC = \angle ABC + \angle ADE$$

$$\Rightarrow$$
 $\angle AD = \angle AE$ [Sides opposite to equal angles are equal]

(i)

Similarly we can prove for Fig (ii) **Proved.**



- **Q.7.** AC and BD are chords of a circle which bisect each other. Prove that (i) AC and BD are diameters, (ii) ABCD is rectangle.
- **Sol. Given :** A circle with chords AB and CD which bisect each other at O.

To Prove: (i) AC and BD are diameters

(ii) ABCD is a rectangle.

Proof : In $\triangle OAB$ and $\triangle OCD$, we have

$$OA = OC$$

[Given]

$$OB = OD$$

[Given]

[Vertically opposite angles]

$$\Rightarrow$$
 $\triangle AOB \cong \angle COD$

[SAS congruence]

$$\Rightarrow$$
 $\angle ABO = \angle CDO$ and $\angle BAO = \angle BCO$

[CPCT]

$$\Rightarrow$$
 AB | | DC

... (i)

Similarly, we can prove BC | | AD

... (ii)

Hence, ABCD is a parallelogram.

But ABCD is a cyclic parallelogram.

: ABCD is a rectangle.

[Proved in Q. 12 of Ex. 10.5]

$$\Rightarrow$$
 $\angle ABC = 90^{\circ} \text{ and } \angle BCD = 90^{\circ}$

⇒ AC is a diameter and BD is a diameter

[Angle in a semicircle is 90°] **Proved.**

Q.8. Bisectors of angles A, B and C of a triangle ABC intersect its circumcircle at D, E and F respectively. Prove that the angles of the triangle DEF are

$$90^{\circ} - \frac{1}{2}A$$
, $90^{\circ} - \frac{1}{2}B$ and $90^{\circ} - \frac{1}{2}C$.

Sol. Given: $\triangle ABC$ and its circumcircle. AD, BE, CF are bisectors of $\angle A$, $\angle B$, $\angle C$ respectively.

Construction: Join DE, EF and FD.

Proof: We know that angles in the same segment are equal.

$$\therefore \qquad \angle 5 = \frac{\angle C}{2} \text{ and } \angle 6 = \frac{\angle B}{2} ...(i)$$

$$\angle 1 = \frac{\angle A}{2}$$
 and $\angle 2 = \frac{\angle C}{2}$..(ii)


$$\angle 4 = \frac{\angle A}{2}$$
 and $\angle 3 = \frac{\angle B}{2}$..(iii)

From (i), we have

$$\angle 5 + \angle 6 = \frac{\angle C}{2} + \frac{\angle B}{2}$$

$$\Rightarrow \angle D = \frac{\angle C}{2} + \frac{\angle B}{2} \qquad ...(iv)$$

 $[\because \angle 5 + \angle 6 = \angle D]$



But $\angle A + \angle B + \angle C = 180^{\circ}$ $\Rightarrow \angle B + \angle C = 180^{\circ} - \angle A$

$$\Rightarrow \frac{\angle B}{2} + \frac{\angle C}{2} = 90^{\circ} - \frac{\angle A}{2}$$

: (iv) becomes,

$$\angle D = 90^{\circ} - \frac{\angle A}{2}$$
.

Similarly, from (ii) and (iii), we can prove that

$$\angle E = 90^{\circ} - \frac{\angle B}{2}$$
 and $\angle F = 90^{\circ} - \frac{\angle C}{2}$ **Proved.**


- **Q.9.** Two congruent circles intersect each other at points A and B. Through A any line segment PAQ is drawn so that P, Q lie on the two circles. Prove that BP = BQ.
- **Sol.** Given: Two congruent circles which intersect at A and B. PAB is a line through A.

To Prove : BP = BQ.

Construction: Join AB.

Proof: AB is a common chord of both the circles.

But the circles are congruent —



$$\Rightarrow$$
 $\angle APB = \angle AQB$ Angles subtended

$$\Rightarrow$$
 BP = BQ [Sides opposite to equal angles are equal] **Proved.**

- **Q.10.** In any triangle ABC, if the angle bisector of $\angle A$ and perpendicular bisector of BC intersect, prove that they intersect on the circumcircle of the triangle ABC.
 - **Sol.** Let angle bisector of $\angle A$ intersect circumcircle of $\triangle ABC$ at D. Join DC and DB.

[Angles in the same segment]

$$\Rightarrow \angle BCD = \angle BAD \frac{1}{2} \angle A$$

[AD is bisector of $\angle A$] ...(i)



From (i) and (ii) $\angle DBC = \angle BCD$

$$\Rightarrow$$
 BD = DC [sides opposite to equal angles are equal]

 \Rightarrow D lies on the perpendicular bisector of BC.

Hence, angle bisector of $\angle A$ and perpendicular bisector of BC intersect on the circumcircle of $\triangle ABC$ **Proved.**

